

- [NCBI](#)
- [Skip to main content](#)
- [Skip to navigation](#)
- [Resources](#)
- [How To](#)
- [About NCBI Accesskeys](#)

[Sign in to NCBI](#)

PMC

[US National Library of Medicine](#)
[National Institutes of Health](#)

Search database

Search term

- [Advanced](#)
- [Journal list](#)
- [Help](#)

- [Journal List](#)
- [HHS Author Manuscripts](#)
- PMC3222922

	HHS Public Access Author manuscript Peer-reviewed and accepted for publication
About author manuscripts	Submit a manuscript

[Int J Cancer](#). Author manuscript; available in PMC 2011 Nov 23.

Published in final edited form as:

[Int J Cancer. 2008 Sep 1; 123\(5\): 1015–1024.](#)

doi: [10.1002/ijc.23588](#)

PMCID: PMC3222922

NIHMSID: NIHMS277965

Activation of PPAR α inhibits IGF-I-mediated growth and survival responses in medulloblastoma cell lines

[Katarzyna Urbanska](#),^{1,2} [Paola Pannizzo](#),¹ [Maja Grabacka](#),³ [Sidney Croul](#),⁴ [Luis Del Valle](#),¹ [Kamel Khalili](#),¹ and [Krzysztof Reiss](#)^{1,*}
[Author information](#) ► [Copyright and License information](#) ►

The publisher's final edited version of this article is available free at [Int J Cancer](#)

See other articles in PMC that [cite](#) the published article.

[Go to:](#)

Abstract

Recent studies suggest a potential role of lipid lowering drugs, fibrates and statins, in anticancer treatment. One candidate for tumor chemoprevention is fenofibrate, which is a potent agonist of peroxisome proliferator activated receptor alpha (PPAR α). Our results demonstrate elevated expression of PPAR α in the nuclei of neoplastic cells in 12 out of 13 cases of medulloblastoma, and of PPAR α in six out of 13 cases. Further analysis demonstrated that aggressive mouse medulloblastoma cells, BsB8, express PPAR α in the absence PPAR γ , and human medulloblastoma cells, D384 and Daoy, express both PPAR α and PPAR γ . Mouse and human cells responded to fenofibrate by a significant increase of PPAR-mediated transcriptional activity, and by a gradual accumulation of cells in G1 and G2/M phase of the cell cycle, leading to the inhibition of cell proliferation and elevated apoptosis. Preincubation of BsB8 cells with fenofibrate attenuated IGF-I-induced IRS-1, Akt, ERKs and GSK3 β phosphorylation, and inhibited clonogenic growth. In Daoy and D384 cells, fenofibrate also inhibited IGF-I-mediated growth responses, and simultaneous delivery of fenofibrate with low dose of the IGF-1R inhibitor, NVP-AEW541, completely abolished their clonogenic growth and survival. These results indicate a strong

the IGF-IR inhibitor, NVP-AEW541, completely abolished their clonogenic growth and survival. These results indicate a strong supportive role of fenofibrate in chemoprevention against IGF-I-induced growth responses in medulloblastoma.

Keywords: PPAR α , fenofibrate, IGF-I, medulloblastoma

Medulloblastomas are highly malignant cerebellar tumors of the childhood, which originate from the external granule layer of the cerebellum and have an inherent tendency to spread in the CNS *via* cerebrospinal fluid. Medulloblastomas are characterized by a large number of genetic and epigenetic aberrations.¹ Among them, overexpression of insulin-like growth factor receptor (IGF-IR), and insulin receptor substrate 1 (IRS-1) are frequently seen in these tumors.¹⁻⁵ The IGF-IR is a membrane-associated tyrosine kinase capable of mediating a variety of biological responses including cell survival and cell proliferation.^{6,7} In the cerebellum, the IGF-IR has been shown in the granule cell layer and in Purkinje cells, and IGF-I protected cultures of cerebellar neurons from low potassium induced apoptosis.^{8,9} In medulloblastoma, the IGF-IR signaling system has been investigated quite intensively.^{4,10} Recent results from our laboratory demonstrate that medulloblastoma cell lines and medulloblastoma biopsies are characterized by an abundant presence of the IGF-IR, and its major signaling molecule, IRS-1.^{2,3} Importantly, we have detected the phosphorylated form of the IGF-IR (active) in the majority of medulloblastoma clinical samples examined.² In addition, growth and survival of medulloblastoma cell lines cultured in suspension was strongly dependent on the presence of exogenous IGF-I.^{2,3} The low-molecular-weight inhibitor of the IGF-IR, NVP-AEW541, is one of the most effective inhibitors of the growth of medulloblastoma cells and induces massive apoptosis in suspension cultures of mouse and human medulloblastoma cell lines.¹¹ We have therefore searched for an additional therapeutic target, which when combined with NVP-AEW541 could enhance the efficacy and decryies toxicity the chemotherapy against medulloblastoma. Previous work in our laboratory indicate that PPARs agonists might serve as the additional agents, because they are known to affect both insulin receptor and IGF-I receptor signaling pathways,^{12,13} and are characterized by relatively low toxicity.^{14,15} PPARs are nuclear receptors, which belong to the superfamily of steroid hormone receptors.¹⁶ Their transcriptional activity after ligand activation is associated with the formation of heterodimers with the retinoid X receptors and binding to the specific DNA sequences.¹⁷ To date, three types of PPARs have been identified: PPAR α , PPAR β and PPAR γ . In the cerebellum all three cellular layers: the molecular, Purkinje cells and granule neurons, express PPARs.^{18,19} Recent reports indicate that activation of PPAR α by fenofibrate attenuates clonogenic growth and migration of melanoma cell lines.^{14,15} Interestingly, fenofibrate has been widely used to lower plasma levels of triglycerides and cholesterol, to improve LDL:HDL ratio, and to prevent the development of arteriosclerosis mainly through regulation of apolipoprotein expression.²⁰ Fenofibrate is also a potent ligand for PPAR α , which has been originally discovered as a regulator of glucose and lipid metabolism, with potential anticancer properties.^{14,15} Since PPARs have not been studied in medulloblastomas, we attempted to ask whether their activation could repress malignant growth of these cerebellar tumors. Our results show elevated levels of PPAR α and PPAR γ in medulloblastoma clinical samples (Fig. 1), and in mouse and human medulloblastoma cell lines (Fig. 2). We have determined also that BsB8 mouse medulloblastoma cells express exclusively PPAR α , and that D384 and Daoy human medulloblastoma cells express both PPAR α and PPAR γ . Mouse and human medulloblastoma cells responded to fenofibrate by a gradual withdrawal from the cell cycle at early time points after the treatment, which resulted in almost complete inhibition of cell proliferation and induction of apoptosis at later time points (after 72 hr). Fenofibrate attenuated IGF-I-induced phosphorylation events, which was accompanied by a severe retardation of their clonogenic growth. Although in human medulloblastoma cell lines fenofibrate was much less effective in triggering apoptosis, simultaneous delivery of fenofibrate with low doses of NVP-AEW541 (0.5 μ M) resulted in a complete inhibition of their clonogenic growth. These results indicate that activation of PPAR α by fenofibrate inhibits cell cycle progression in the first 48 hr, and after prolonged treatment (over 72 hr) it can induce apoptosis. Because of the low toxicity of fenofibrate,^{14,15} it could be considered as a supplementary treatment against IGF-I-mediated growth responses in medulloblastoma.




FIGURE 1

Immunohistochemical detection of PPARs in medulloblastoma clinical samples. Evaluation of PPAR α and PPAR γ protein levels in medulloblastomas in comparison to normal cerebellar tissue. (a-c) View of the normal cerebellum where the ...




FIGURE 2

Peroxisome proliferator-activated receptors (PPARs) in medulloblastoma cell lines. (a) Western blot showing PPAR α and PPAR γ in two human (D384, Daoy) and one mouse (BsB8) medulloblastoma cell lines. Brown adipose tissue was used as a positive ...

[Go to:](#)

Material and methods

Immunohistochemistry

A total of 13 cases of human medulloblastoma were acquired from the pathology archives of the University Health Network, University of Toronto, Ontario, Canada. The formalin-fixed, paraffin-embedded samples were sectioned at 4 microns in thickness, stained with Hematoxylin and Eosin and histologically classified according to the World Health Organization Classification of Tumors of the Nervous System. Immunohistochemistry was performed using the avidin–biotin–peroxidase complex system, according to the manufacturer's instructions (Vectastain Elite ABC Peroxidase Kit; Vector Laboratories, Burlingame, CA). Briefly, sections were deparaffinized in xylene and rehydrated through descending alcohols to water. For nonenzymatic antigen retrieval sections were heated in 0.01 M sodium citrate buffer (pH 6.0) to 95°C for 40 min and allowed to cool for 20 min at room temperature. To quench endogenous peroxidase, slides were then rinsed with PBS and incubated in MeOH/3% H₂O₂ for 20 min. Sections were then washed with PBS and blocked in PBS/0.1% BSA containing 5% normal horse or goat serum for 2 hr at room temperature, and incubated overnight at room temperature with primary antibodies. The primary antibodies used in our study include: a mouse monoclonal anti-PPAR α antibody (1:100 dilution; Chemicon, Temecula, CA), a rabbit polyclonal anti-PPAR γ (E-8, 1:300 dilution; Santa Cruz Biotechnology, Santa Cruz, CA), a mouse monoclonal anti-class III β -tubulin (TUJ1, 1:500 dilution; Covance, Berkeley, CA) and a mouse monoclonal cocktail for different molecular weight neurofilaments (SMI-312, 1:2000 dilution; Sternbeger Monoclonals, Lutherville, MA). Secondary antibodies used were biotin-conjugated horse anti-mouse and goat anti-rabbit IgGs (Vector). Sections were developed with a diaminobenzidine substrate (Sigma, St. Louis, MO), counterstained with hematoxylin and mounted with Permount.

Cell culture conditions

The following cell lines were used in our study: a mouse medulloblastoma, BsB8, isolated from a cerebellar tumor of a transgenic mouse expressing the early genome of human polyomavirus JC^{3,21}; and two human medulloblastoma cell lines, D384 Med and Daoy. D384 are metastatic medulloblastomas isolated from peritoneal ascites of a child with medulloblastoma,²² and Daoy are derived from a tumor in the posterior fossa of a 4-years-old boy (ATCC no. HTB186). BsB8 cells are very aggressive, capable of forming large tumors in subcutaneous tissue, as well as intracranial tumors after stereotactic injection of a small number of cells in to the brain, and have the ability of spreading into cerebellum.^{1,23} BsB8 and Daoy were maintained as monolayer cultures in Dulbecco's modified Eagle medium (DMEM) (GIBCO-BRL, Grand Island, NY) containing 10% fetal bovine serum (FBS), at 37°C in a 9.6% CO₂ atmosphere. D384 were cultured in suspension, in the Eagle's MEM supplemented with nonessential amino acids (GIBCO-BRL), 2 mM l-glutamine, 1 mM sodium pyruvate, and 20% FBS. The cells were made partially quiescent by 48 hr incubation in serum-free medium (SFM) (DMEM supplemented with 0.1% bovine serum albumin). Cellular growth and signaling responses were tested by stimulating serum starved cells with 50 ng/ml of recombinant IGF-I (GIBCO-BRL).

Luciferase assay

The PPARs transcriptional activity in BsB8 and D384 cells was determined by utilizing the JsTkpGL3 reporter plasmid, which contains the luciferase gene driven by PPAR responsive element (PPRE), which consists of three copies of the J site from apo-All gene promoter.²⁴ The activation of PPAR elements was evaluated by a dual-Firefly/Renilla luciferase reporter system (Promega, Madison, WI), using Femtomaster FB12 chemiluminometer (Zylux Corp., Oak Ridge, TN).

Western blot analysis

To evaluate phosphorylation levels of the selected IGF-IR signaling molecules, semiquiescent monolayer cultures were stimulated with IGF-I and total protein extracts collected at the indicated time points. The cells were lysed for 5 min on ice with 400 μ l of lysis buffer A [50 mM HEPES, pH 7.5; 150 mM NaCl; 1.5 mM MgCl₂; 1 mM EGTA; 10% glycerol; 1% Triton X-100; 1 μ M phenylmethylsulfonyl fluoride (PMSF); 0.2 mM Na-ortho-vanadate; 10 μ g/ml aprotinin]. Total protein extracts (50 μ g) were separated on a 4–15% gradient SDS-PAGE (BioRad, Hercules, CA) and transferred to nitrocellulose filters. The following primary anti-phospho antibodies were utilized: anti-pY612IRS-1 rabbit polyclonal (BioSource, Camarillo, CA); anti-pS473Akt, anti-pT202/Y204Erk1/2, and anti-pS21/9GSK3 α / β (Cell Signaling Technology, Beverly, MA). In addition, anti-PPAR α mouse monoclonal antibody (Chemicon) and anti-PPAR γ rabbit polyclonal antibody (Santa Cruz Biotechnology) were utilized. To monitor loading conditions we used anti-IRS-1 (Upstate USA, Charlottesville, VA), anti-GSK3 β , anti-Akt, anti-Erks antibodies (Cell Signaling, Danvers, MA) and anti-Grb-2 antibodies.

Cell cycle analysis and apoptosis

Aliquots of cells, 1×10^6 /ml, were fixed in 70% ethanol for 30 min at 4°C. Cells were centrifuged at 1,600 rpm and the resulting pellets were resuspended in 1 ml of freshly prepared propidium iodide/RNaseA solution. Cell cycle distribution was analyzed with GuavaEasy Cyte system by using Guava CytoSoft cell cycle program according to the manufacturer recommendations (Guava Technologies, Hayward, CA). Based on differences in PI intensity of fluorescence, cells in G1, S, G2/M and subG1 were separated and counted. Each cell cycle experiment was repeated at least three times. Cells in subG1 are characterized by DNA content, which is below the DNA content in G1 (single copy of the genome), and are considered as a mixture of cells dying by apoptosis and/or necrosis. To monitor apoptosis, cells were additionally labeled by in situ terminal dUTP nick-end labeling (TUNEL) assay. We have utilized the Guava TUNEL kit and the Guava TUNEL EasyCyte program to determine the percentage of medulloblastoma cells undergoing apoptosis at different time points after fenofibrate treatment.

Cell growth in monolayer

BsB8 cells were seeded at a concentration of 2×10^4 cells/35 mm dish, in DMEM containing 10% FBS and 50ng/ml IGF-I. 24 hr after

plating cells were either treated with 25 μ M fenofibrate, or were left untreated. At 48 and 72 hr time points, the cells were collected by trypsinization and counted in a Bright-line Hemocytometer in the presence of Trypan blue. T0 is the cell number determined 16 hr after the initial plating of the cells, which reflects plating efficiency.

Clonogenic assay

Cells were plated at 1×10^3 on 6-well culture plates (Costar, Corning, NY) in DMEM supplemented with 10% FBS and 50 ng/ml of IGF-I. After attachment, cells were treated with fenofibrate, NVP-AEW541 (kindly provided by Novartis Pharma, Basel, Switzerland), or with combination of fenofibrate and NVP-AEW541 at the indicated concentration. Clonogenic potential was evaluated 10 days after the treatment by fixing and staining corresponding cell cultures with 0.25% Brilliant Blue in methanol, which was followed by macroscopic count of the clones.

Colony formation in soft agar

To assess anchorage-independent growth of D384, cells were plated at 1×10^4 per 35 mm/dish in DMEM containing 10% FBS, 50 ng/ml of IGF-I and 0.2% agarose, with 0.4% agarose underlay. The growth medium was additionally supplemented with 25 μ M fenofibrate or with a combination of 25 μ M fenofibrate and 0.5 μ M NVP-AEW541. Clones larger than 125 μ m in diameter were scored after continuous growth of the cells for 2 weeks.

[Go to:](#)

Results

Detection of PPARs in medulloblastoma clinical samples

The results in [Figure 1](#) demonstrate nuclear expression of PPAR α and PPAR γ in a significant number of tumor cells from two representative medulloblastoma biopsies ([Figs. 1e and 1f](#)). In normal human cerebellum ([Fig. 1a–1c](#)), PPAR α was detected only in proximity to Purkinje cells, where a population of neurofilament positive Basket cells ([Fig. 1a](#); arrow) was strongly positive for PPAR α ([Fig. 1b](#); arrows). In contrast, immunolabeling with anti-PPAR γ antibody of normal human cerebellum (including molecular layer, granular layer, Purkinje cells and Basket cells) was negative ([Fig. 1c](#)). We have analyzed a total of 13 medulloblastoma clinical samples, and detected PPAR α in 92%, and PPAR γ in 46% of all medulloblastoma cases examined.

Fenofibrate -mediated activation of PPAR α in medulloblastoma cell lines

We have previously reported strong antitumoral properties of a synthetic PPAR α agonist, fenofibrate, which in melanoma cell lines attenuated constitutive phosphorylation of Akt.¹⁵ To investigate whether fenofibrate can affect IGF-I-mediated growth responses in medulloblastomas, we selected the aggressive mouse medulloblastoma cell line, BsB8, and two human medulloblastoma cell lines Daoy and D384, which are strongly responsive to the IGF-I stimulation.^{3,4,11} The western blot analysis depicted in [Fig. 2a](#) shows that BsB8, Daoy and D384 cells express PPAR α protein at levels comparable to the control brown adipose tissue. In contrast to human medulloblastoma cell lines, which express both PPAR α and PPAR γ , BsB8 cells were negative for PPAR γ . Our finding was additionally confirmed in JCV T-antigen transgenic mice ([Fig. 2b](#)), which develop cerebellar primitive neuroectodermal tumors, from which BsB8 were originally isolated.^{3,21} The luciferase based transcriptional activity assay demonstrated that the fenofibrate treatment resulted in three-fold and four-fold increase in the activity of PPRES in BsB8 and D384 cells, respectively ([Fig. 2c](#)), further confirming PPAR α -mediated transcriptional activity in mouse and human medulloblastoma cell lines.

PPAR α -mediated inhibition of growth responses

We then evaluated the monolayer growth of BsB8 cells, the condition in which medulloblastoma cells are quite resistant to the treatment against the IGF-IR function.¹¹ The cells were plated in 10%FBS in the presence (Feno) or absence (control) of 25 μ M fenofibrate.¹⁵ The cell number was determined at time zero (T0; 16 hr after initial plating), and at 48 and 72 hr after the fenofibrate treatment. The results in [Figure 3a](#) show that in comparison to BsB8 cells growing in the presence of 10%FBS, fenofibrate-treated cells demonstrated a 4-fold lower rate of cell proliferation. This growth inhibition was accompanied by morphological changes ([Fig. 3b](#)), including accumulation of perinuclear structures within first 24 hr after the treatment (arrows), which likely represents increased peroxisome proliferation;²⁵ the appearance of spindle shaped large nondividing cells, between 48 and 72 hr (arrowheads), and apoptotic cells in later time points ([Fig. 4b](#); subG1 cell population).




FIGURE 3

Inhibition of growth responses of BsB8 cells after PPAR α activation. (a) BsB8 mouse medulloblastomas were plated in the presence of 10% FBS in the presence (Feno) or absence (control) of 25 μ M fenofibrate. The cell number was determined at time zero (T0; 16 hr after initial plating), and at 48 and 72 hr after the fenofibrate treatment. The results in [Figure 3a](#) show that in comparison to BsB8 cells growing in the presence of 10%FBS, fenofibrate-treated cells demonstrated a 4-fold lower rate of cell proliferation. This growth inhibition was accompanied by morphological changes ([Fig. 3b](#)), including accumulation of perinuclear structures within first 24 hr after the treatment (arrows), which likely represents increased peroxisome proliferation;²⁵ the appearance of spindle shaped large nondividing cells, between 48 and 72 hr (arrowheads), and apoptotic cells in later time points ([Fig. 4b](#); subG1 cell population).

of 10% FBS at $2 \times 10^7/35$ mm dish. Attached cells were treated with $25 \mu\text{M}$ fenofibrate (F-eno) or were ...




FIGURE 4

Effects of fenofibrate on IGF-I signaling pathways. (a) After 24 hr incubation in serum-free medium (SFM), quiescent BSB8 were cultured in the presence or absence of fenofibrate for additional 24 hr. Subsequently, both cell groups were treated with IGF-I ...

Effects of fenofibrate on IGF-I signaling pathways

Since fenofibrate modulates insulin-mediated cellular responses, and the IGF-IR signaling system is highly active in medulloblastomas,¹ we asked whether the observed growth inhibitory effects of fenofibrate are associated with the inhibition of IGF-IR signaling pathway(s). After 24 hr incubation in SFM, semiquiescent Bsb8 cells were cultured in the presence or absence of $25 \mu\text{M}$ fenofibrate for an additional 24 hrs. Subsequently, the cells were treated with IGF-I (50 ng/ml) for 30 min, 6 hr, or were left without the IGF-I treatment (SFM). The results in [Figure 4a](#) demonstrate a strong increase in the phosphorylation of IRS-1, Akt, Erks1/2 and GSK-3 β during first 30 min after IGF-stimulation, and an expected desensitization of the signal at the 6 hr time point. Preincubation of the cells with $25 \mu\text{M}$ fenofibrate attenuated IGF-I-mediated phosphorylation events ([Fig. 4a](#)). Cell cycle analysis shown in [Fig. 4b](#) revealed accumulation of cells in G2/M phase, and decrease in the S phase of the cell cycle at the 48 hr time point, which was followed by the accumulation of cells in subG1 phase 72 hr after the treatment. This subG1 cells fraction (23.7%) was most likely associated with the loss of DNA content in cells undergoing apoptosis. Indeed, TUNEL assay demonstrated that 31% \pm 6 of Bsb8 cells underwent apoptosis 72 hr after the fenofibrate treatment. These inhibitory effects of fenofibrate were almost completely abolished by GW-9662, which at $10 \mu\text{M}$ concentration is considered as a strong PPAR α and PPAR γ antagonist.¹⁵ In a similar manner, pre-incubation of Bsb8 cells with $25 \mu\text{M}$ fenofibrate inhibited IGF-I-stimulated clonogenic growth ([Fig. 4c](#)).

Since the human medulloblastoma cell lines, Daoy and D384, express both PPAR α and PPAR γ , we asked whether fenofibrate could inhibit their growth responses to IGF-I in a similar manner to Bsb8 cells. The results in [Figure 5](#) illustrate that fenofibrate partially downregulated IGF-I-mediated phosphorylation of IRS-1, Akt, ERKs and Gsk-3 β , which was associated with dose dependent inhibition of the clonogenic growth of Daoy cells by fenofibrate ([Fig. 5b](#)). Although fenofibrate-mediated inhibition of growth was quite effective ([Fig. 5b](#)), expected apoptotic cell death was less apparent than in Bsb8 cells (compare [Figs. 4b](#) and [and5c](#)). Also, in contrast to Bsb8 cells in which G2/M arrest predominated, fenofibrate induced partial G1 arrest in Daoy cells, which was accompanied by only slightly elevated apoptosis (from 2% in control samples to 4% in fenofibrate treated samples) ([Fig. 5c](#)).




FIGURE 5

Effects of fenofibrate on IGF-I-mediated cell signaling responses in Daoy cells. (a) After 48 hr incubation in serum-free medium (SFM), quiescent Daoy were cultured in the presence or absence of fenofibrate for additional 24 hr. The cells were subsequently ...

We attempted next to sensitize human medulloblastoma cell lines to fenofibrate by utilizing low concentrations of a specific IGF-IR inhibitor, NVP-AEW541 (Novartis). We have previously shown that medulloblastomas are sensitive to $1 \mu\text{M}$ NVP-AEW541 treatment when kept in suspension cultures, and were much more resistant to the same treatment when cultured in monolayer.¹¹ The results depicted in [Fig. 6](#) indicate that simultaneous treatment of Daoy and D384 cells with fenofibrate ($25 \mu\text{M}$) and NVP-AEW541 ($0.5 \mu\text{M}$) caused almost complete inhibition of the clonogenic growth of Daoy cells ([Fig. 6a](#)), and inhibited colony formation of D384 cells in soft agar ([Fig. 6b](#)). We have obtained 97.3% inhibition of the colony formation in soft agar in the presence of $0.5 \mu\text{M}$ NVP AEW541 (ref.) and $25 \mu\text{M}$ fenofibrate ([Fig. 6b](#)). Note also that D384 cells were partially resistant to $1 \mu\text{M}$ NVP AEW541 in soft agar assay.¹¹ Results in [Figure 6C](#) revealed that cell cycle distribution of D384 cell growing in suspension was quite different from the cell cycle obtained from Bsb8 and Daoy cells, which grew as monolayer cultures (compare [Figs. 4b](#), [5c](#), and [and6c](#)). The major difference was associated with relatively high level of subG1 fraction of the cells in optimal growth conditions (8.8%). Fenofibrate treatment increased the accumulation of cells in subG1 phase from 8.8% to 32.8% during first 24 hr of the treatment, and this high rate of apoptosis was also detected in 72 hr time point. At this late time point the percentage of cells in S and G2/M phase of the cell cycle also decreased, indicating that both cell death and attenuation of cell cycle progression contribute to the inhibitory action of fenofibrate on D384 cells.






FIGURE 6

Growth inhibition of human medulloblastoma cell lines by fenofibrate and NVP-AEW541. (a) Clonogenic growth of Daoy cells. The cells were plated at clonal density (1×10^3 per 35 mm dish), in the presence of 10% FBS, FBS + IGF-I (50 ng/ml) or in ...

[Go to:](#)

Discussion

The presented results demonstrate for the first time the nuclear presence of PPAR α and PPAR γ proteins in medulloblastoma cell lines and in medulloblastoma biopsies. Mouse and human tumor cells responded to fenofibrate treatment with a significant increase of PPAR-mediated transcriptional activity, and an accumulation of cells in G2/M phase (BsB8 mouse cells) and G1 phase (Daoy and D384 human cells) of the cell cycle. Pre-incubation of BsB8 cells with 25 μ M fenofibrate for 24 hr attenuated IGF-I-induced IRS-1, Akt, ERKs and GSK3 β phosphorylation, which resulted in a severe retardation of their clonogenic growth. In Daoy and D384 human cells, preincubation with fenofibrate partially inhibited IGF-I-mediated phosphorylation and growth responses; and together with a low dose of the IGF-IR inhibitor, NVP-AEW541, fenofibrate completely inhibited clonogenic growth and colony formation in soft agar. This indicates an important role of fenofibrate in chemoprevention against IGF-I-induced growth responses in medulloblastoma.

Growth and survival of medulloblastoma cells in vitro strongly depends on the activation of IGF-IR.^{11,26} In medulloblastoma clinical samples the IGF-IR and its major signaling molecule, IRS-1, are strongly upregulated and active (tyrosine phosphorylated).² Different molecular and pharmacological manipulations directed against the IGF-IR were effective in attenuating medulloblastoma growth and survival in vitro,^{3,11} and in experimental animals.^{1,23} Our results demonstrate that 24 hr pre-incubation of mouse and human medulloblastoma cell lines with fenofibrate inhibited IGF-I-mediated phosphorylation events (Figs. 4a and 5a). This result could be seen as a surprise since fenofibrate has been found to increase insulin-mediated signaling responses including enhanced glucose uptake, mitochondrial glucose oxidation and reduced adiposity,¹³ and is frequently used to counteract metabolic abnormalities associated with insulin resistance in type-2 diabetes.^{27–29} However, fenofibrate treatment inhibited VEGF- and bFGF-induced endothelial cells migration, which was accompanied by a decrease in Akt phosphorylation.^{30,31} In melanoma cells, fenofibrate attenuated both cell invasiveness and clonogenic growth by inhibiting constitutively active in these tumors Akt.^{14,15} In contrast, some authors report a rapid but transient increase of Erk1/2 and Akt phosphorylation caused by PPAR α and PPAR γ agonists.^{32–34} In those studies, changes in the phosphorylation were detected shortly after fenofibrate treatment (in minutes), and therefore, were not dependent on the canonical activity of PPAR α as a transcription factor.^{34,35} In our experiments, fenofibrate induced PPARs transcriptional activity and the biological consequences of its action accumulated over time, leading to cell cycle arrest, and to apoptotic cell death in later time points after the treatment (Fig. 4b). These long lasting cytostatic and cytotoxic responses to fenofibrate and the expression of PPAR α in medulloblastoma clinical samples (Fig. 1) indicate a potential therapeutic application of fenofibrate against medulloblastoma.

An alternative concept of PPAR α antitumoral action is associated with cancer cell energy metabolism. This notion has been initiated by Otto Warburg, who indicated a distinctive dependence of tumor cell metabolism from glycolysis, even when there is sufficient amount of oxygen available for much more effective oxidative phosphorylation.^{36,37} Only recently, it has been established that the inclination of tumor cells for glycolysis is mainly driven by mitochondrial dysfunction.^{38,39} PPAR α , which is a transcriptional activator of fatty acid β -oxidation machinery (e.g. acyl-CoA oxidase, acyl-CoA synthetase, carnitine palmitoyl transferase, fatty acid binding protein, fatty acid transporter), can switch energy metabolism towards fatty acid degradation, and decrease glucose uptake by repressing glucose transporter GLUT4.^{40,41} Interestingly, PPAR α acts as a direct sensor for fatty acids, which are considered as natural ligands for this nuclear receptor.^{42,43} According to fatty acid–glucose cycle paradigm, increased rate of fatty acid and ketone bodies oxidation forces the decline in glucose utilization through the inhibition of glycolytic enzymes.^{44,45} This concept was supported by the results of animal studies, showing that during fasting activated PPAR α can divert energy metabolism from the glucose to fatty acid utilization as a primary source of energy. In addition, loss-of-function mutations of genes encoding the Krebs cycle enzymes (such as succinate dehydrogenase and fumarate hydratase) are observed in uterine leiomyomas, renal carcinomas, paragangliomas and pheochromocytomas.⁴⁶ Therefore, glycolysis-promoting metabolism of the cancer cell could relieve the selection pressure and support growth and survival of the cells with defective mitochondrial system. Note, however, that such cells could be brought to the verge of metabolic catastrophe in the condition of limited glucose availability, or when the oxidative metabolism is forced pharmacologically. This opens an opportunity for the use of PPAR α ligands, including fenofibrate, as they should be selectively toxic for cancer cells and neutral for normal cells.

In mouse medulloblastoma cells, the treatment with 25 μ M fenofibrate was sufficient to stop cell proliferation, and to induce massive apoptosis (Fig. 4b). The concentration of fenofibrate was comparable to plasma concentrations of fenofibric acid, an active metabolite of fenofibrate, detected in patients during standard hyperlipidemia treatment (300 mg regular or 250 mg slow release capsules daily).

In such patients, the plateau phase plasma concentration of fenofibric acid is approximately 10–12 μ g/ml (28–33 μ M).⁴⁷ Therefore, 25 μ M fenofibrate holds promise for low toxicity systemic chemotherapy against medulloblastoma in comparison to current regimens many of which are highly toxic.^{48,49} In human medulloblastoma cells, Daoy and D384, which express PPAR α and PPAR γ , 25 μ M fenofibrate attenuated IGF-I-mediated signaling events including IRS-1, Akt, ERKs and GSK-3 β phosphorylation (Fig. 5a), and

triggered cell cycle arrest (Fig. 5b). However, there was induction of apoptosis than in BsB8 cells (Fig. 4b). Although increasing fenofibrate concentration to 50 μ M induced apoptosis in Daoy cells (not shown), this higher concentration could be systemically toxic. We therefore tested the effects of 25 μ M fenofibrate in combination with low doses (up to 0.5 μ M) of the IGF-IR inhibitor NVP-AEW541. This low-molecular-weight inhibitor of the IGF-IR kinase activity has been previously shown to limit the growth and survival of medulloblastoma cell lines in anchorage-independent culture conditions.¹¹ Interestingly, medulloblastoma cells in monolayer cultures were quite resistant to the treatment with 1 μ M NVP-AEW541.¹¹ In our study, lower dose of NVP-AEW541 (0.5 μ M) used in combination with 25 μ M fenofibrate completely inhibited clonogenic growth of Daoy cells (Fig. 6a), and efficiently inhibited colony formation of D384 cells in soft agar (Fig. 6b).

Although the mechanism by which fenofibrate attenuates IGF-I signaling responses is still under investigation, it has been recently reported that fenofibrate increases plasma membrane rigidity in a manner similar to the elevated cholesterol content in cell membranes.⁵⁰ In that report, fenofibrate did not change the membrane content of cholesterol, but increased plasma membrane rigidity, altering activities of integral membrane proteins such as sarco (endo)plasmic reticulum Ca^{2+} -ATPase and β -secretase-mediated cleavage of APP.⁵⁰ Further experiments are required to determine whether similar fenofibrate-mediated changes in the fluidity of plasma membrane are indeed responsible for the attenuation of ligand-induced clusterization of the IGF-IR molecules—a critical step in autophosphorylation of the receptor molecules and the initiation of growth promoting signaling cascades.

[Go to:](#)

Acknowledgements

We thank Dr. Shohreh Amini for her editorial help.

Grant sponsor: NIH; Grant numbers: RO1CA095518-01 and PO1 NS36466-06.

[Go to:](#)

References

1. Reiss K. Insulin-like growth factor-I receptor—a potential therapeutic target in medulloblastomas. *Expert Opin Ther Targets*. 2002;6:539–544. [[PubMed](#)]
2. Del Valle L, Enam S, Lassak A, Wang JY, Croul S, Khalili K, Reiss K. Insulin-like growth factor I receptor activity in human medulloblastomas. *Clin Cancer Res*. 2002;8:1822–1830. [[PubMed](#)]
3. Wang JY, Del Valle L, Gordon J, Rubini M, Romano G, Croul S, Peruzzi F, Khalili K, Reiss K. Activation of the IGF-IR system contributes to malignant growth of human and mouse medulloblastomas. *Oncogene*. 2001;20:3857–3868. [[PubMed](#)]
4. Patti R, Reddy CD, Georger B, Grotzer MA, Raghunath M, Sutton LN, Phillips PC. Autocrine secreted insulin-like growth factor-I stimulates MAP kinase-dependent mitogenic effects in human primitive neuroectodermal tumor/medulloblastoma. *Int J Oncol*. 2000;16:577–584. [[PubMed](#)]
5. Rao G, Pedone CA, Valle LD, Reiss K, Holland EC, Fufts DW. Sonic hedgehog and insulin-like growth factor signaling synergize to induce medulloblastoma formation from nestin-expressing neural progenitors in mice. *Oncogene*. 2004;23:6156–6162. [[PubMed](#)]
6. Baserga R, Sell C, Porcu P, Rubini M. The role of the IGF-I receptor in the growth and transformation of mammalian cells. *Cell Prolif*. 1994;27:63–71. [[PubMed](#)]
7. Baserga R, Peruzzi F, Reiss K. The IGF-1 receptor in cancer biology. *Int J Cancer*. 2003;107:873–877. [[PubMed](#)]
8. D'Mello SR, Galli C, Ciotti T, Calissano P. Induction of apoptosis in cerebellar granule neurons by low potassium: inhibition of death by insulin-like growth factor I and cAMP. *Proc Natl Acad Sci USA*. 1993;90:10989–10993. [[PMC free article](#)] [[PubMed](#)]
9. Cui H, Meng Y, Bulleit RF. Inhibition of glycogen synthase kinase 3beta activity regulates proliferation of cultured cerebellar granule cells. *Brain Res Dev Brain Res*. 1998;111:177–188. [[PubMed](#)]
10. Kurihara M, Tokunaga Y, Ochi A, Kawaguchi T, Tsutsumi K, Shigematsu K, Niwa M, Mori K. [Expression of insulin-like growth factor I receptors in human brain tumors: comparison with epidermal growth factor receptor by using quantitative autoradiography] *No To Shinkei*. 1989;41:719–725. [[PubMed](#)]
11. Urbanska K, Trojanek J, Del Valle L, Eldeen MB, Hofmann F, Garcia-Echeverria C, Khalili K, Reiss K. Inhibition of IGF-I receptor in anchorage-independence attenuates GSK-3beta constitutive phosphorylation and compromises growth and survival of medulloblastoma cell lines. *Oncogene*. 2007;26:2308–2317. [[PubMed](#)]
12. Lecka-Czernik B, Ackert-Bicknell C, Adamo ML, Marmolejos V, Churchill GA, Shockley KR, Reid IR, Grey A, Rosen CJ. Activation of peroxisome proliferator-activated receptor gamma (PPARgamma) by rosiglitazone suppresses components of the insulin-like growth factor regulatory system in vitro and in vivo. *Endocrinology*. 2007;148:903–911. [[PMC free article](#)] [[PubMed](#)]
13. Guerre-Millo M, Gervois P, Raspe E, Madsen L, Poulain P, Derudas B, Herbert JM, Winegar DA, Willson TM, Fruchart JC, Berge RK, Staels B. Peroxisome proliferator-activated receptor alpha activators improve insulin sensitivity and reduce adiposity. *J Biol Chem*. 2000;275:16638–16642. [[PubMed](#)]
14. Grabacka M, Placha W, Plonka PM, Pajak S, Urbanska K, Laidler P, Slominski A. Inhibition of melanoma metastases by fenofibrate. *Arch Dermatol Res*. 2004;296:54–58. [[PubMed](#)]
15. Grabacka M, Plonka PM, Urbanska K, Reiss K. Peroxisome proliferator-activated receptor alpha activation decreases metastatic potential of melanoma cells in vitro via down-regulation of Akt. *Clin Cancer Res*. 2006;12:3028–3036. [[PubMed](#)]
16. Issemann I, Green S. Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature*. 1990;347:645–650. [[PubMed](#)]
17. Kliewer SA, Umehara K, Noonan DJ, Heyman RA, Evans RM. Convergence of 9-cis retinoic acid and peroxisome proliferator signalling pathways through heterodimer formation of their receptors. *Nature*. 1992;358:771–774. [[PubMed](#)]
18. Braissant O, Fufelle F, Scotto C, Dauca M, Wahli W. Differential expression of peroxisome proliferator-activated receptors (PPARs): tissue distribution of PPAR-alpha, -beta, and -gamma in the adult rat. *Endocrinology*. 1996;137:354–366. [[PubMed](#)]
19. Kainu T, Wikstrom AC, Gustafsson JA, Pelto-Huikko M. Localization of the peroxisome proliferator-activated receptor in the brain



19. Kanda T, Wikstrom AC, Gustafsson OA, Petroniuk M. Localization of the peroxisome proliferator-activated receptor in the brain. *Neuroreport*. 1994;5:2481–2485. [[PubMed](#)]
20. Staels B, Van Tol A, Fruchart JC, Auwerx J. Effects of hypolipidemic drugs on the expression of genes involved in high density lipoprotein metabolism in the rat. *Isr J Med Sci*. 1996;32:490–498. [[PubMed](#)]
21. Krynska B, Otte J, Franks R, Khalili K, Croul S. Human ubiquitous JCV(CY) T-antigen gene induces brain tumors in experimental animals. *Oncogene*. 1999;18:39–46. [[PubMed](#)]
22. He XM, Wikstrand CJ, Friedman HS, Bigner SH, Pleasure S, Trojanowski JQ, Bigner DD. Differentiation characteristics of newly established medulloblastoma cell lines (D384 Med, D425 Med, and D458 Med) and their transplantable xenografts. *Lab Invest*. 1991;64:833–843. [[PubMed](#)]
23. Wang JY, Del Valle L, Peruzzi F, Trojanek J, Giordano A, Khalili K, Reiss K. Polyomaviruses and cancer—interplay between viral proteins and signal transduction pathways. *J Exp Clin Cancer Res*. 2004;23:373–383. [[PubMed](#)]
24. Vu-Dac N, Schoonjans K, Kosykh V, Dallongeville J, Fruchart JC, Staels B, Auwerx J. Fibrates increase human apolipoprotein A-II expression through activation of the peroxisome proliferator-activated receptor. *J Clin Invest*. 1995;96:741–750. [[PMC free article](#)] [[PubMed](#)]
25. Lores Arnaiz S, Travacio M, Monserrat AJ, Cutrin JC, Llesuy S, Boveris A. Chemiluminescence and antioxidant levels during peroxisome proliferation by fenofibrate. *Biochim Biophys Acta*. 1997;1360:222–228. [[PubMed](#)]
26. Wang H, Zeng ZC, Bui TA, Sonoda E, Takata M, Takeda S, Iliakis G. Efficient rejoining of radiation-induced DNA double-strand breaks in vertebrate cells deficient in genes of the RAD52 epistasis group. *Oncogene*. 2001;20:2212–2224. [[PubMed](#)]
27. Bajaj M, Suraamornkul S, Hardies LJ, Glass L, Musi N, Defronzo RA. Effects of peroxisome proliferator-activated receptor (PPAR)-alpha and PPAR-gamma agonists on glucose and lipid metabolism in patients with type 2 diabetes mellitus. *Diabetologia*. 2007;8:1723–1731. [[PubMed](#)]
28. Tan CE, Chew LS, Tai ES, Chio LF, Lim HS, Loh LM, Shepherd J. Benefits of micronised Fenofibrate in type 2 diabetes mellitus subjects with good glycemic control. *Atherosclerosis*. 2001;154:469–474. [[PubMed](#)]
29. Wu TJ, Ou HY, Chou CW, Hsiao SH, Lin CY, Kao PC. Decrease in inflammatory cardiovascular risk markers in hyperlipidemic diabetic patients treated with fenofibrate. *Ann Clin Lab Sci*. 2007;37:158–166. [[PubMed](#)]
30. Goetze S, Eilers F, Bungenstock A, Kintscher U, Stawowy P, Blaschke F, Graf K, Law RE, Fleck E, Grafe M. PPAR activators inhibit endothelial cell migration by targeting Akt. *Biochem Biophys Res Commun*. 2002;293:1431–1437. [[PubMed](#)]
31. Varet J, Vincent L, Mirshahi P, Pille JV, Legrand E, Opolon P, Mishal Z, Soria J, Li H, Soria C. Fenofibrate inhibits angiogenesis in vitro and in vivo. *Cell Mol Life Sci*. 2003;60:810–819. [[PubMed](#)]
32. Baek SJ, Wilson LC, Hsi LC, Eling TE. Troglitazone, a peroxisome proliferator-activated receptor gamma (PPAR gamma) ligand, selectively induces the early growth response-1 gene independently of PPAR gamma. A novel mechanism for its anti-tumorigenic activity. *J Biol Chem*. 2003;278:5845–5853. [[PubMed](#)]
33. Gardner OS, Dewar BJ, Earp HS, Samet JM, Graves LM. Dependence of peroxisome proliferator-activated receptor ligand-induced mitogen-activated protein kinase signaling on epidermal growth factor receptor transactivation. *J Biol Chem*. 2003;278:46261–46269. [[PubMed](#)]
34. Takeda K, Ichiki T, Tokunou T, Iino N, Takeshita A. 15-Deoxy-delta 12,14-prostaglandin J2 and thiazolidinediones activate the MEK/ERK pathway through phosphatidylinositol 3-kinase in vascular smooth muscle cells. *J Biol Chem*. 2001;276:48950–48955. [[PubMed](#)]
35. Pauley CJ, Ledwith BJ, Kaplanski C. Peroxisome proliferators activate growth regulatory pathways largely via peroxisome proliferator-activated receptor alpha-independent mechanisms. *Cell Signal*. 2002;14:351–358. [[PubMed](#)]
36. Warburg O. On respiratory impairment in cancer cells. *Science*. 1956;124:269–270. [[PubMed](#)]
37. Warburg O. On the origin of cancer cells. *Science*. 1956;123:309–314. [[PubMed](#)]
38. Degenhardt K, Mathew R, Beaudoin B, Bray K, Anderson D, Chen G, Mukherjee C, Shi Y, Gelinas C, Fan Y, Nelson DA, Jin S, et al. Autophagy promotes tumor cell survival and restricts necrosis, inflammation, and tumorigenesis. *Cancer Cell*. 2006;10:51–64. [[PMC free article](#)] [[PubMed](#)]
39. Shaw RJ. Glucose metabolism and cancer. *Curr Opin Cell Biol*. 2006;18:598–608. [[PubMed](#)]
40. Ahmed W, Ziouzenkova O, Brown J, Devchand P, Francis S, Kadakia M, Kanda T, Orasanu G, Shirlach M, Zandbergen F, Plutzky J. PPARs and their metabolic modulation: new mechanisms for transcriptional regulation? *J Intern Med*. 2007;262:184–198. [[PubMed](#)]
41. Finck BN, Kelly DP. Peroxisome proliferator-activated receptor alpha (PPARalpha) signaling in the gene regulatory control of energy metabolism in the normal and diseased heart. *J Mol Cell Cardiol*. 2002;34:1249–1257. [[PubMed](#)]
42. Forman BM, Chen J, Evans RM. Hypolipidemic drugs, polyunsaturated fatty acids, and eicosanoids are ligands for peroxisome proliferator-activated receptors alpha and delta. *Proc Natl Acad Sci USA*. 1997;94:4312–4317. [[PMC free article](#)] [[PubMed](#)]
43. Xu HE, Lambert MH, Montana VG, Parks DJ, Blanchard SG, Brown PJ, Sternbach DD, Lehmann JM, Wisely GB, Willson TM, Kliewer SA, Milburn MV. Molecular recognition of fatty acids by peroxisome proliferator-activated receptors. *Mol Cell*. 1999;3:397–403. [[PubMed](#)]
44. Randle PJ. Regulatory interactions between lipids and carbohydrates: the glucose fatty acid cycle after 35 years. *Diabetes Metab Rev*. 1998;14:263–283. [[PubMed](#)]
45. Wolfe RR. Metabolic interactions between glucose and fatty acids in humans. *Am J Clin Nutr*. 1998;67:519S–526S. [[PubMed](#)]
46. Brandon M, Baldi P, Wallace DC. Mitochondrial mutations in cancer. *Oncogene*. 2006;25:4647–4662. [[PubMed](#)]
47. Doser K, Guserle R, Nitsche V, Arnold G. Comparative steady state study with 2 fenofibrate 250 mg slow release capsules. An example of bioequivalence assessment with a highly variable drug. *Int J Clin Pharmacol Ther*. 1996;34:345–348. [[PubMed](#)]
48. Corden BJ, Strauss LC, Killmond T, Carson BS, Wharam MD, Kumar AJ, Piantadosi S, Robb PA, Phillips PC. Cisplatin, ara-C and etoposide (PAE) in the treatment of recurrent childhood brain tumors. *J Neurooncol*. 1991;11:57–63. [[PubMed](#)]
49. Knight KR, Kraemer DF, Neuwelt EA. Ototoxicity in children receiving platinum chemotherapy: underestimating a commonly occurring toxicity that may influence academic and social development. *J Clin Oncol*. 2005;23:8588–8596. [[PubMed](#)]
50. Gamerding M, Clement AB, Behl C. Cholesterol-like effects of selective COX inhibitors and fibrates on cellular membranes and amyloid- β production. *Mol Pharmacol*. 2007;72:141–151. [[PubMed](#)]

Formats:

- Article

- [PubReader](#)
- [ePub \(beta\)](#)
- [PDF \(5.6M\)](#)
- [Citation](#)

Share

-  [Facebook](#)
-  [Twitter](#)
-  [Google+](#)

Save items

[Add to Favorites](#)[View more options](#)

loading

1

Similar articles in PubMed

- [Inhibition of IGF-I receptor in anchorage-independence attenuates GSK-3beta constitutive phosphorylation and compromises growth and survival of medulloblastoma cell lines.](#)[Oncogene. 2007]
- [ROS accumulation and IGF-IR inhibition contribute to fenofibrate/PPARalpha -mediated inhibition of glioma cell motility in vitro.](#)[Mol Cancer. 2010]
- [Fenofibrate suppresses growth of the human hepatocellular carcinoma cell via PPARalpha-independent mechanisms.](#)[Eur J Cell Biol. 2011]
- [Activation of PPAR alpha by fenofibrate inhibits apoptosis in vascular adventitial fibroblasts partly through SIRT1-mediated deacetylation of FoxO1.](#)[Exp Cell Res. 2015]
- [The insulin-like growth factor-I receptor kinase inhibitor NVP-ADW742 sensitizes medulloblastoma to the effects of chemotherapy.](#)[Oncol Rep. 2011]

[See reviews...](#)[See all...](#)

Cited by other articles in PMC

- [PPAR-gamma Agonists As Antineoplastic Agents in Cancers with Dysregulated IGF Axis](#)[Frontiers in Endocrinology. 2017]
- [Stimulating effect of normal-dosing of fibrates on cell proliferation: word of warning](#)[Lipids in Health and Disease. ...]
- [Fenofibrate Suppresses Oral Tumorigenesis via Reprogramming Metabolic Processes: Potential Drug Repurposing for Oral Cancer](#)[International Journal of Biolo...]
- [Metformin stimulates IGFBP-2 gene expression through PPARalpha in diabetic states](#)[Scientific Reports. 2016]
- [PPARalpha modulates gene expression profiles of mitochondrial energy metabolism in oral tumorigenesis](#)[BioMedicine. 2016]

[See all...](#)

Links

- [Compound](#)
- [MedGen](#)
- [PubMed](#)
- [Substance](#)
- [Taxonomy](#)

Recent Activity

[Clear](#)[Turn Off](#)[Turn On](#)

- [Activation of PPARalpha inhibits IGF-I-mediated growth and survival responses in med...](#)
Activation of PPARalpha inhibits IGF-I-mediated growth and survival responses in medulloblastoma cell lines
NIHPA Author Manuscripts. 2008 Sep 1; 123(5)1015
- [Central nervous system concentrations of cyclooxygenase-2 inhibitors in humans.](#)
Central nervous system concentrations of cyclooxygenase-2 inhibitors in humans.
Anesthesiology. 2005 Feb;102(2):409-15.
PubMed

- [COX-2 drives metastatic breast cells from brain lesions into the cerebrospinal f...](#)
COX-2 drives metastatic breast cells from brain lesions into the cerebrospinal fluid and systemic circulation. Cancer Res. 2014 May 1;74(9):2385-90. doi: 10.1158/0008-5472.CAN-13-2660. Epub 2014 Mar 10. PubMed
- [Practice advisory: Etanercept for poststroke disability](#)
Practice advisory: Etanercept for poststroke disability Neurology. 2016 Jun 7; 86(23):2208
- [Comment on: "Perispinal Etanercept for Post-Stroke Neurological and Cognitive Dy...](#)
Comment on: "Perispinal Etanercept for Post-Stroke Neurological and Cognitive Dysfunction: Scientific Rationale and Current Evidence". CNS Drugs. 2014 Dec;28(12):1205-6. doi: 10.1007/s40263-014-0211-1. PubMed

Your browsing activity is empty.

Activity recording is turned off.

[Turn recording back on](#)

[See more...](#)

- [Activation of the IGF-IR system contributes to malignant growth of human and mouse medulloblastomas.](#)[Oncogene. 2001]
Wang JY, Del Valle L, Gordon J, Rubini M, Romano G, Croul S, Peruzzi F, Khalili K, Reiss K
Oncogene. 2001 Jun 28; 20(29):3857-68.
- [Human ubiquitous JCV\(CY\) T-antigen gene induces brain tumors in experimental animals.](#)[Oncogene. 1999]
Krynska B, Otte J, Franks R, Khalili K, Croul S
Oncogene. 1999 Jan 7; 18(1):39-46.
- [Differentiation characteristics of newly established medulloblastoma cell lines \(D384 Med, D425 Med, and D458 Med\) and their transplantable xenografts.](#)[Lab Invest. 1991]
He XM, Wikstrand CJ, Friedman HS, Bigner SH, Pleasure S, Trojanowski JQ, Bigner DD
Lab Invest. 1991 Jun; 64(6):833-43.
- [Insulin-like growth factor-I receptor - a potential therapeutic target in medulloblastomas.](#)[Expert Opin Ther Targets. 2002]
Reiss K
Expert Opin Ther Targets. 2002 Oct; 6(5):539-44.
- [Review Polyomaviruses and cancer--interplay between viral proteins and signal transduction pathways.](#)[J Exp Clin Cancer Res. 2004]
Wang JY, Del Valle L, Peruzzi F, Trojanek J, Giordano A, Khalili K, Reiss K
J Exp Clin Cancer Res. 2004 Sep; 23(3):373-83.
- [Fibrates increase human apolipoprotein A-II expression through activation of the peroxisome proliferator-activated receptor.](#)[J Clin Invest. 1995]
Vu-Dac N, Schoonjans K, Kosykh V, Dallongeville J, Fruchart JC, Staels B, Auwerx J
J Clin Invest. 1995 Aug; 96(2):741-50.
- [Peroxisome proliferator-activated receptor alpha activation decreases metastatic potential of melanoma cells in vitro via down-regulation of Akt.](#)[Clin Cancer Res. 2006]
Grabacka M, Plonka PM, Urbanska K, Reiss K
Clin Cancer Res. 2006 May 15; 12(10):3028-36.
- [Activation of the IGF-IR system contributes to malignant growth of human and mouse medulloblastomas.](#)[Oncogene. 2001]
Wang JY, Del Valle L, Gordon J, Rubini M, Romano G, Croul S, Peruzzi F, Khalili K, Reiss K
Oncogene. 2001 Jun 28; 20(29):3857-68.
- [Autocrine secreted insulin-like growth factor-I stimulates MAP kinase-dependent mitogenic effects in human primitive neuroectodermal tumor/medulloblastoma.](#)[Int J Oncol. 2000]
Patti R, Reddy CD, Georger B, Grotzer MA, Raghunath M, Sutton LN, Phillips PC
Int J Oncol. 2000 Mar; 16(3):577-84.
- [Inhibition of IGF-I receptor in anchorage-independence attenuates GSK-3beta constitutive phosphorylation and compromises growth and survival of medulloblastoma cell lines.](#)[Oncogene. 2007]
Urbanska K, Trojanek J, Del Valle L, Eldeen MB, Hofmann F, Garcia-Echeverria C, Khalili K, Reiss K
Oncogene. 2007 Jul 12; 26(28):3998-4008.

- Oncogene. 2007 Apr 5; 26(16):2308-17.
- [Human ubiquitous JCV\(CY\) T-antigen gene induces brain tumors in experimental animals.](#)[Oncogene. 1999]
Krynska B, Otte J, Franks R, Khalili K, Croul S
Oncogene. 1999 Jan 7; 18(1):39-46.
 - [Inhibition of IGF-I receptor in anchorage-independence attenuates GSK-3beta constitutive phosphorylation and compromises growth and survival of medulloblastoma cell lines.](#)[Oncogene. 2007]
Urbanska K, Trojanek J, Del Valle L, Eldeen MB, Hofmann F, Garcia-Echeverria C, Khalili K, Reiss K
Oncogene. 2007 Apr 5; 26(16):2308-17.
 - [Peroxisome proliferator-activated receptor alpha activation decreases metastatic potential of melanoma cells in vitro via down-regulation of Akt.](#)[Clin Cancer Res. 2006]
Grabacka M, Plonka PM, Urbanska K, Reiss K
Clin Cancer Res. 2006 May 15; 12(10):3028-36.
 - [Chemiluminescence and antioxidant levels during peroxisome proliferation by fenofibrate.](#)[Biochim Biophys Acta. 1997]
Lores Arnaiz S, Travacio M, Monserrat AJ, Cutrín JC, Llesuy S, Boveris A
Biochim Biophys Acta. 1997 May 24; 1360(3):222-8.
 - [Insulin-like growth factor-I receptor - a potential therapeutic target in medulloblastomas.](#)[Expert Opin Ther Targets. 2002]
Reiss K
Expert Opin Ther Targets. 2002 Oct; 6(5):539-44.
 - [Peroxisome proliferator-activated receptor alpha activation decreases metastatic potential of melanoma cells in vitro via down-regulation of Akt.](#)[Clin Cancer Res. 2006]
Grabacka M, Plonka PM, Urbanska K, Reiss K
Clin Cancer Res. 2006 May 15; 12(10):3028-36.
 - [Inhibition of IGF-I receptor in anchorage-independence attenuates GSK-3beta constitutive phosphorylation and compromises growth and survival of medulloblastoma cell lines.](#)[Oncogene. 2007]
Urbanska K, Trojanek J, Del Valle L, Eldeen MB, Hofmann F, Garcia-Echeverria C, Khalili K, Reiss K
Oncogene. 2007 Apr 5; 26(16):2308-17.
 - [Inhibition of IGF-I receptor in anchorage-independence attenuates GSK-3beta constitutive phosphorylation and compromises growth and survival of medulloblastoma cell lines.](#)[Oncogene. 2007]
Urbanska K, Trojanek J, Del Valle L, Eldeen MB, Hofmann F, Garcia-Echeverria C, Khalili K, Reiss K
Oncogene. 2007 Apr 5; 26(16):2308-17.
 - [Efficient rejoining of radiation-induced DNA double-strand breaks in vertebrate cells deficient in genes of the RAD52 epistasis group.](#)[Oncogene. 2001]
Wang H, Zeng ZC, Bui TA, Sonoda E, Takata M, Takeda S, Iliakis G
Oncogene. 2001 Apr 26; 20(18):2212-24.
 - [Insulin-like growth factor I receptor activity in human medulloblastomas.](#)[Clin Cancer Res. 2002]
Del Valle L, Enam S, Lassak A, Wang JY, Croul S, Khalili K, Reiss K
Clin Cancer Res. 2002 Jun; 8(6):1822-30.
 - [Activation of the IGF-IR system contributes to malignant growth of human and mouse medulloblastomas.](#)[Oncogene. 2001]
Wang JY, Del Valle L, Gordon J, Rubini M, Romano G, Croul S, Peruzzi F, Khalili K, Reiss K
Oncogene. 2001 Jun 28; 20(29):3857-68.
 - [Insulin-like growth factor-I receptor - a potential therapeutic target in medulloblastomas.](#)[Expert Opin Ther Targets. 2002]
Reiss K
Expert Opin Ther Targets. 2002 Oct; 6(5):539-44.
 - [Review Polyomaviruses and cancer--interplay between viral proteins and signal transduction pathways.](#)[J Exp Clin Cancer Res. 2004]
Wang JY, Del Valle L, Peruzzi F, Trojanek J, Giordano A, Khalili K, Reiss K
J Exp Clin Cancer Res. 2004 Sep; 23(3):373-83.
 - [Peroxisome proliferator-activated receptor alpha activators improve insulin sensitivity and reduce adiposity.](#)[J Biol Chem. 2000]
Guerre-Millo M, Gervois P, Raspé E, Madsen L, Poulain P, Derudas B, Herbert JM, Winegar DA, Willson TM, Fruchart JC, Berge RK, Staels B
J Biol Chem. 2000 Jun 2; 275(22):16638-42.

- [Effects of peroxisome proliferator-activated receptor \(PPAR\)-alpha and PPAR-gamma agonists on glucose and lipid metabolism in patients with type 2 diabetes mellitus.](#)[Diabetologia. 2007]
Bajaj M, Suraamornkul S, Hardies LJ, Glass L, Musi N, DeFronzo RA
Diabetologia. 2007 Aug; 50(8):1723-31.
- [Decrease in inflammatory cardiovascular risk markers in hyperlipidemic diabetic patients treated with fenofibrate.](#)[Ann Clin Lab Sci. 2007]
Wu TJ, Ou HY, Chou CW, Hsiao SH, Lin CY, Kao PC
Ann Clin Lab Sci. 2007 Spring; 37(2):158-66.
- [PPAR activators inhibit endothelial cell migration by targeting Akt.](#)[Biochem Biophys Res Commun. 2002]
Goetze S, Eilers F, Bungenstock A, Kintscher U, Stawowy P, Blaschke F, Graf K, Law RE, Fleck E, Gräfe M
Biochem Biophys Res Commun. 2002 May 24; 293(5):1431-7.
- [Fenofibrate inhibits angiogenesis in vitro and in vivo.](#)[Cell Mol Life Sci. 2003]
Varet J, Vincent L, Mirshahi P, Pille JV, Legrand E, Opolon P, Mishal Z, Soria J, Li H, Soria C
Cell Mol Life Sci. 2003 Apr; 60(4):810-9.

[See more ...](#)

- [On respiratory impairment in cancer cells.](#)[Science. 1956]
WARBURG O
Science. 1956 Aug 10; 124(3215):269-70.
- [On the origin of cancer cells.](#)[Science. 1956]
WARBURG O
Science. 1956 Feb 24; 123(3191):309-14.
- [Autophagy promotes tumor cell survival and restricts necrosis, inflammation, and tumorigenesis.](#)[Cancer Cell. 2006]
Degenhardt K, Mathew R, Beaudoin B, Bray K, Anderson D, Chen G, Mukherjee C, Shi Y, Gélinas C, Fan Y, Nelson DA, Jin S, White E
Cancer Cell. 2006 Jul; 10(1):51-64.
- [Review Glucose metabolism and cancer.](#)[Curr Opin Cell Biol. 2006]
Shaw RJ
Curr Opin Cell Biol. 2006 Dec; 18(6):598-608.
- [Review PPARs and their metabolic modulation: new mechanisms for transcriptional regulation?](#)[J Intern Med. 2007]
Ahmed W, Ziouzenkova O, Brown J, Devchand P, Francis S, Kadakia M, Kanda T, Orasanu G, Sharlach M, Zandbergen F, Plutzky J
J Intern Med. 2007 Aug; 262(2):184-98.
- [Review Peroxisome proliferator-activated receptor alpha \(PPARalpha\) signaling in the gene regulatory control of energy metabolism in the normal and diseased heart.](#)[J Mol Cell Cardiol. 2002]
Finck BN, Kelly DP
J Mol Cell Cardiol. 2002 Oct; 34(10):1249-57.
- [Hypolipidemic drugs, polyunsaturated fatty acids, and eicosanoids are ligands for peroxisome proliferator-activated receptors alpha and delta.](#)[Proc Natl Acad Sci U S A. 1997]
Forman BM, Chen J, Evans RM
Proc Natl Acad Sci U S A. 1997 Apr 29; 94(9):4312-7.
- [Molecular recognition of fatty acids by peroxisome proliferator-activated receptors.](#)[Mol Cell. 1999]
Xu HE, Lambert MH, Montana VG, Parks DJ, Blanchard SG, Brown PJ, Sternbach DD, Lehmann JM, Wisely GB, Willson TM, Kliewer SA, Milburn MV
Mol Cell. 1999 Mar; 3(3):397-403.
- [Review Regulatory interactions between lipids and carbohydrates: the glucose fatty acid cycle after 35 years.](#)[Diabetes Metab Rev. 1998]
Randle PJ
Diabetes Metab Rev. 1998 Dec; 14(4):263-83.
- [Review Metabolic interactions between glucose and fatty acids in humans.](#)[Am J Clin Nutr. 1998]
Wolfe RR
Am J Clin Nutr. 1998 Mar; 67(3 Suppl):519S-526S.
- [Review Mitochondrial mutations in cancer.](#)[Oncogene. 2006]
Brandon M, Baldi P, Wallace DC
Oncogene. 2006 Aug 7; 25(34):4647-62.
- [Comparative steady state study with 2 fenofibrate 250 mg slow release capsules. An example of bioequivalence assessment with a highly variable drug.](#)[Int J Clin Pharmacol Ther. 1996]

Doser K, Guserle R, Nitsche V, Arnold G
Int J Clin Pharmacol Ther. 1996 Aug; 34(8):345-8.

- [Cisplatin, ara-C and etoposide \(PAE\) in the treatment of recurrent childhood brain tumors.](#)[J Neurooncol. 1991]
Corden BJ, Strauss LC, Killmond T, Carson BS, Wharam MD, Kumar AJ, Piantadosi S, Robb PA, Phillips PC
J Neurooncol. 1991 Aug; 11(1):57-63.
- [Ototoxicity in children receiving platinum chemotherapy: underestimating a commonly occurring toxicity that may influence academic and social development.](#)[J Clin Oncol. 2005]
Knight KR, Kraemer DF, Neuwelt EA
J Clin Oncol. 2005 Dec 1; 23(34):8588-96.
- [Inhibition of IGF-I receptor in anchorage-independence attenuates GSK-3beta constitutive phosphorylation and compromises growth and survival of medulloblastoma cell lines.](#)[Oncogene. 2007]
Urbanska K, Trojanek J, Del Valle L, Eldeen MB, Hofmann F, Garcia-Echeverria C, Khalili K, Reiss K
Oncogene. 2007 Apr 5; 26(16):2308-17.
- [Cholesterol-like effects of selective cyclooxygenase inhibitors and fibrates on cellular membranes and amyloid-beta production.](#)[Mol Pharmacol. 2007]
Gamerding M, Clement AB, Behl C
Mol Pharmacol. 2007 Jul; 72(1):141-51.

[Support Center](#) [Support Center](#)

External link. Please review our [privacy policy](#).

[NLM](#)

[NIH](#)

[DHHS](#)

[USA.gov](#)

[National Center for Biotechnology Information, U.S. National Library of Medicine](#) 8600 Rockville Pike, Bethesda MD, 20894 USA

[Policies and Guidelines](#) | [Contact](#)

•

